

CELLULAR THERAPIES – DELIVERY OF THEM: LEARNING FROM CAR-T CELLS

Dr Tobias Menne MD, PhD, MRCP, FRCPath

Clinical Director for Research

Consultant Haematologist

CAR-T cell lead

The Newcastle upon Tyne Hospitals



DISCLOSURES DR MENNE

- Travel grants: Amgen, Jazz, Pfizer, Bayer, Kyowa Kirin, Celgene/BMS, Kite/Gilead, Janssen, Takeda
- Honoraria for advisory board meetings: Kite/Gilead, Amgen, Novartis, Pfizer, Celgene/BMS, Daiichi Sankyo, Atara, Roche, Janssen
- Honoraria for lectures: Kite/Gilead, Takeda, Janssen, Roche, Servier, Novartis, Celgene/BMS, Pfizer, Incyte
- Research funding: Janssen, AstraZeneca, Novartis



UK EXPERIENCE


NICE recommendations

- Oct 18 Axicel for 3L+ HGBCL
- Nov 18 Tisagen for R/R-B-ALL up to age 25 years
- Nov 18 Tisagen for 3L+ HGBCL
- Feb 21 Brexu-cel for R/R MCL
- June 23 Brexucel for RR B-ALL 26+ years
- June 23 Axicel for 2nd line HGBCL
- Mar 25 Lisocel for 2nd line HGBCL
- Summer 25 Obecel for RR B-ALL (Expected)

- **Note**


So far no Myeloma CAR-T cell therapy has been approved by NICE due to cost implications


VERY COMPLEX
PROCESS
TO DELIVER
CAR-T CELLS



TOPICS TO COVER

- Apheresis phase
- Manufacturing CAR-T cell phase
- Lymphodepletion and CAR-T treatment phase
- Post CAR-T therapy phase






APHERESIS PHASE


- Identify patients
 - Does the patient meet the NICE approved indication
 - ECOG performance status 0 or I
 - Disease that is stable enough to allow collection of T-cells and manufacture of CAR-T cells
 - “Adequate organ function”
- Discuss at national CAR-T panel
- Apherese as quickly as possible

MANUFACTURING
CAR-T CELL
PHASE


- Typically takes 3-4 weeks from apheresis to start lymphodepletion
- Often Bridging therapy is given
 - Chemotherapy
 - Radiotherapy
- Risk of progression with deteriorating ECOG performance score
- Low chance of CAR-T production failure or OOS - Out of Specification product




Number at risk									
None	40	26	21	14	12	10	9	4	2
Steroids	29	13	8	7	6	6	6	5	0
RT	54	39	33	26	14	10	8	0	0
Systemic	166	90	65	40	27	15	10	4	1
CMT	11	8	5	4	4	3	1	1	1



Number at risk									
SD/PD	115	58	38	28	19	13	9	2	1
CR/PR	104	74	62	39	24	14	10	3	1



Number at risk									
None	40	37	34	23	18	15	13	5	3
Steroids	29	23	14	12	10	7	7	6	0
RT	54	50	42	30	17	13	8	0	0
Systemic	166	132	111	71	42	24	14	6	2
CMT	11	9	8	6	5	4	1	1	1



Number at risk									
SD/PD	115	86	66	49	31	20	12	4	2
CR/PR	104	94	86	52	30	19	11	3	1

EFFECTIVE BRIDGING CAN IMPROVE CAR-T OUTCOMES

Multivariate analysis of baseline factors

Factors affecting response to BT	Responder/N	OR (95% CI)	P-value
RBP bridging			
No	34/134	1.00	.010
Yes	35/83	2.21 (1.21-4.05)	
Response last line			
SD/PD	45/165	1.00	.023
CR/PR	24/52	2.16 (1.11-4.22)	
Bulky disease			
No	55/149	1.00	.045
Yes	14/68	0.49 (0.25-0.98)	
Factors affecting response to CAR-T	Events/N	HR (95% CI)	P-value
LDH at LD			
≤2ULN	74/141	1.00	.001
>2ULN	34/41	2.06 (1.34-3.16)	
Extra nodal sites			
<3	91/160	1.00	.001
≥3	17/22	2.51 (1.46-4.32)	
BT response			
SD/PD	68/100	1.00	.012
CR/PR	40/82	0.58 (0.38-0.89)	

LYMPHODEPLETION
AND
CAR-T TREATMENT
PHASE

- Lymphodepletion phase 5-6 days followed by CAR-T infusion
- CAR-T treatment phase typically 28 days with most patients being an inpatient for 10-14 days post CAR-T infusion
- High chance of developing CRS and ICANS
- TRM rates of 3-5%
- Response assessment at 28-30 days post CAR-T cells important predictor for PFS

LARGE B-CELL LYMPHOMA PATIENTS TREATED WITH CD19 CAR T IN THE UK OVER TIME

2nd line Axi-cel
UK RWD data


N=195 patients
Median FU 9.1 months

ORR: 88%
CMR: 61%

Patients – Axi-cel infused
9-month PFS: 47.6%
9-month OS: 77%

Patients – not infused
9-month OS: 19.7


PFS of infused patients by treatment Era



No at risk	0	6	12	18	24	30	36
Era 2	407	232	171	108	18	0	0
Era 1	171	66	55	53	51	30	4


12-month OS (95% CI)	All patients:	53.1% (49.4% - 56.8%)
	Era 2:	59.6% (55.1% - 64.1%)
	Era 1:	40.3% (34.0% - 46.6%)

OS (ITT) from CAR-T approval date by Era



No at risk	0	6	12	18	24	30	36
Era 2	490	349	260	175	73	0	0
Era 1	236	143	96	76	70	51	16


PFS by Treatment Era and Product Choice



No at risk	0	6	12	18	24	30	36
Axi-cel Era 2	338	199	149	88	18	0	0
Tisa-cel Era 2	67	34	24	20	1	0	0
Axi-cel Era 1	120	50	42	40	38	24	2
Tisa-cel Era 1	49	15	12	12	12	8	1


12-month PFS (95% CI)	All patients:	44.3% (40.2% - 48.4%)
	Era 2, Age <70:	49.9% (44.2% - 55.6%)
	Era 2, Age >70:	49.2% (39.0% - 59.4%)
	Era 1, Age <70:	33.3% (25.9% - 40.8%)
	Era 1, Age >70:	16.7% (0.00% - 34.0%)

PFS by Treatment Era and Age



No at risk	0	6	12	18	24	30	36
<70y Era 2	312	182	132	85	17	0	0
70+ Era 2	93	52	39	23	2	0	0
<70y Era 1	152	60	51	48	47	29	3
70+ Era 1	17	5	3	3	3	3	0


EARLY FDG-PET RESPONSE PREDICTS CAR-T FAILURE IN LARGE B-CELL LYMPHOMA



MANAGEMENT OF TOXICITIES

EARLY AND LATE

TIMING OF LIFE THREATENING COMPLICATIONS FOLLOWING CAR-T CELLS INFUSION




ACUTE CAR-T TOXICITY OUTCOMES IN THE UK OVER TIME



	Era 1 2019 <i>n</i> (available)	%	Era 2 2020-2022 <i>n</i> (available)	%	<i>p</i> Value	2 nd line- UK 5/2023- 3/2024 N=195 %
Of those who received CAR T						
Axi-cel infused	121 (171)	71	339 (407)	83	<0.001	Axi-cel infused 91
CRS grade 3+	16 (171)	9	16 (405)	4	0.01	CRS grade 3+ 4
ICANS grade 3+	27 (171)	16	63 (405)	16	0.94	ICANS Grade 3+ 17
Tocilizumab use	104 (171)	61	141 (185)	76	0.002	Tocilizumab use 90
Corticosteroid use	59 (171)	35	84 (184)	46	0.03	Corticosteroid use 60
ICU admission	55 (171)	32	79 (405)	20	0.001	ICU admission 20
G3+ neutropenia at 3 months	26 (133)	20	16 (92)	17	0.68	Anakinra use 17
G3+ thrombocytopenia at 3 months	25 (133)	19	18 (92)	20	0.88	G3+ ICAHT at 1 month 29
NRM at 12 months	10 (167)	6	30 (402)	7.5	0.53	NRM at 9 months 5.4

POST CAR-T
THERAPY
PHASE

- From Day 28 onwards
- Shared care model is standard of care
- Ongoing infection, cytopenias and immunosuppression remains significant issue
- Many CAR-T centres have established nurse-led “late toxicity” clinics



NON-RELAPSE MORTALITY AFTER STANDARD OF CARE AXI-CEL IN 3L INFECTIOUS COMPLICATIONS




US Lymphoma CART Consortium - RWD dataset with 5 year follow up
 275 patients treated with Axicel in 3L+

Death due to infections (majority respiratory): total 21/275 (7.7%)

- Age <60 years 4/132 (3%)
- Age >60 years 17/143 (11.8%)


B-CELL RECOVERY AFTER CAR-T CELLS (AXICEL)



Severe presentations and high mortality from SARS-CoV-2 in patients undergoing chimeric antigen receptor (CAR-T) therapy: a UK NCCP analysis


Kathleen P. L. Cheok^{a,b}, Amy A. Kirkwood^c, Tobias Menne^d, Eleni Tholouli^e, Sridhar Chaganti^f, Amrith Mathew^f, Ben Uttenthal^g, James Russell^g, David Irvine^h, Rod Johnsonⁱ, Emma Nicholson^j, Jessica Bazin^j, William Townsend^b, Andrea Kuhn^k , Maeve O'Reilly^b, Robin Sanderson^k, Amit Patel^l and Claire Roddie^{a,b}

- Post CAR-T infusion
 - high mortality rate if patient develops severe COVID19
 - 12/14 died (86%)



SHARED CARE MODEL

2L CAR-T TREATMENT PATHWAY





CONCLUSIONS

- Really interesting times for CAR-T cell therapies
- More indications becoming available and CAR-T cells move into earlier therapy lines
- Best sequencing between CAR-T cell and bispecific therapies needs to be established
- Allogenic CAR-T cell therapies are looking promising